

***Psyttoma* gen. n. (Hymenoptera, Braconidae, Opiinae) from Shandong and Hubei (China), with a key to the species**

Xi-Ying Li^{1,†}, Cornelis van Achterberg^{2,‡}, Ji-Cai Tan^{3,§}

1 College of Bio-Safety Science and Technology, Hunan Agriculture University, Changsha 410128, China **2** Department of Terrestrial Zoology, Naturalis Biodiversity Center, Postbus 9517, 2300 RA Leiden, The Netherlands

† [urn:lsid:zoobank.org:author:A2CF381E-FB87-404D-899F-98408B8FE556](https://zoobank.org/A2CF381E-FB87-404D-899F-98408B8FE556)

‡ [urn:lsid:zoobank.org:author:D6374CF4-8F07-4EA8-8C55-9335FD19CECD](https://zoobank.org/D6374CF4-8F07-4EA8-8C55-9335FD19CECD)

§ [urn:lsid:zoobank.org:author:5C77DC9F-FD52-4510-99B3-6CD87350F7E8](https://zoobank.org/5C77DC9F-FD52-4510-99B3-6CD87350F7E8)

Corresponding author: Cornelis van Achterberg (Cees.vanAchterberg@naturalis.nl)

Academic editor: Gavin Broad | Received 25 March 2012 | Accepted 30 August 2012 | Published 15 October 2012

[urn:lsid:zoobank.org:pub:1AC4D606-321C-4EDC-A7A1-4C5161C8B31F](https://zoobank.org/pub/1AC4D606-321C-4EDC-A7A1-4C5161C8B31F)

Citation: Li X-Y, Achterberg C van, Tan J-C (2012) *Psyttoma* gen. n. (Hymenoptera, Braconidae, Opiinae) from Shandong and Hubei (China), with a key to the species. Journal of Hymenoptera Research 29: 73–81. doi: 10.3897/JHR.29.3123

Abstract

Psyttoma gen. n. (type species: *Opius latilabris* Chen & Weng, 2005) from Shandong and Hubei (China) is described and illustrated (Hymenoptera: Braconidae: Opiinae). *Psyttoma latilabris* (Chen & Weng, 2005) and *P. wachsmanni* (Szépligeti, 1898) are new combinations.

Keywords

Braconidae, Opiinae, *Psyttoma*, *Psytalia*, new genus, Palaearctic, Oriental, China, Shandong, Hubei

Introduction

The large subfamily Opiinae (Braconidae), with 1,975 valid species according to Yu et al. (2009), is a common group containing generally small (2–5 mm) parasitoid wasps of mainly mining or fruit-infesting dipterous larvae. It has a worldwide distribution and the world fauna has been reviewed by Fischer (1972, 1977, 1986, 1987). Currently about 35 genera are used; Wharton (1988, 1997), van Achterberg (1997b, 2004a,

2004b), van Achterberg and Salvo (1997) and van Achterberg and Chen (2004) published updates or some additions for the existing keys to the genera of the Opiinae, but the number of genera and the limits of several genera are still matters of discussion.

The first author collected some Opiinae for her revision of the Opiinae of Hunan in her home town in the province of Shandong (Palearctic China). One of the species resembles members of the genus *Psytalia* Walker, 1860, but does not fit well in this genus. Also DNA analysis in the DNA lab of the Naturalis Biodiversity Center (Leiden) showed that also the molecular data disagree with inclusion in the genera *Psytalia*, *Opius* or *Phaenodrotoma*. A combined analysis of the 16S and 28S sequences of about 50 mainly Oriental Opiine taxa from China and Malaysia was made and the results corroborated the view that the new genus has a separate position among the Opiine genera. The results of this study will be published elsewhere (Li et al., in prep.). Therefore, we describe the species in a new genus below. The DNA sequences are deposited in GenBank and have been compared with unpublished sequences of *Psytalia* species from Malaysia and with published sequences (Gimeno et al., 1997). Later it was discovered that the mutilated type of *Opius latilabris* Chen & Weng, 2005, from Hubei (Oriental China) belongs to this species.

Opiinae are solitary koinobiont endoparasitoids of larvae of cyclorrhaphous Diptera, but oviposition may take place in the egg of the host (ovo-larval parasitoid). They may play an important role in the control of such dipterous pests as fruit-infesting Tephritidae and mining Agromyzidae. The parasitoid larva completes its final development when the host larva has made its puparium and the adult emerges from this puparium.

Material and methods

The specimens were directly killed in 70% alcohol during collection. The redescribed specimens are deposited in the Naturalis collection (RMNH) at Leiden. For identification of the subfamily Opiinae, see van Achterberg (1990, 1993, and 1997a); for identification of the genera, see Chen and Weng (2005), Wharton (1988, 1997), van Achterberg (1997b, 2004a); for references to the Opiinae, see Yu et al. (2009); and for the terminology used in this paper, see van Achterberg (1988, 1993). Measurements are taken as indicated by van Achterberg (1988).

Taxonomy

Psytoma van Achterberg & Li, gen. n.

[urn:lsid:zoobank.org:act:29FD6792-00E8-441F-9C72-E97458B50D3A](http://zoobank.org/act:29FD6792-00E8-441F-9C72-E97458B50D3A)

<http://species-id.net/wiki/Psyttoma>

Figures 1–13

Etymology. The name is a combination of the generic names *Psytalia* Walker and *Phaenodrotoma* Foerster, 1862, because it combines features of both genera. Gender: feminine.




Figure 1. *Psyttoma latilabris* (Chen & Weng), female from Anqiu. Habitus lateral.

Type species. *Opius latilabris* Chen & Weng, 2005.

Diagnosis. Hypopygium of female distinctly acute apically and about 0.3 times as long as metasoma; labrum slanted backwards, leaving a large space below clypeus (Fig. 9); veins 2-SR+R and SC+R1 of hind wing of male strongly widened (Fig. 13); precoxal sulcus smooth; hind femur 2–3 times as long as wide (Fig. 11); hind tibia without carinula basally; scutellum convex medially and depressed medio-posteriorly, distinctly protruding above level of mesoscutum (Fig. 5); propodeum without median carina or areola; second metasomal segment 1.2 times longer than third tergite (Fig. 4); spiracle of second tergite distinctly removed from apex of first tergite (Fig. 4).

Notes. The genus will run in the key to world genera of Opiinae by Wharton (1997) to the genera *Psyttalia* Walker and *Opius* Wesmael (part with normal mandible; = *Phaedrotoma*). Because of the sharp pointed hypopygium it seems closer to *Psyttalia*, but it does not fit in any one of the genera and differs in having a medio-posterior depression of the mesoscutum, the second metasomal tergite longer than the third tergite, the precoxal sulcus smooth, the scutellum protruding dorsally, the hind wing narrowed and vein m-cu of the hind wing present as a faintly pigmented trace. The new genus can be separated from both genera as follows:

- 1 Scutellum distinctly protruding above level of mesoscutum (Fig. 5); hypopygium of ♀ distinctly acute apically and about 0.3 times as long as metasoma (Fig. 7) and hind wing comparatively narrow (Figs 12, 13); hind femur very robust, 2-3 times as long as wide (Fig. 11); labrum slanted backwards,

- leaving a large space below clypeus (Fig. 9); medio-anterior veins of hind wing of male strongly widened (Fig. 13) ***Psyttoma* gen. n.**
- Scutellum at level of mesoscutum; hypopygium of ♀ variable, if distinctly acute apically and about 0.3 times as long as metasoma then hind wing moderately wide and hind femur slender, 4–5 times as long as wide; labrum normal, without large space below clypeus; medio-anterior veins of hind wing of male narrow ***Psyttalia* Walker and *Phaedrotoma* Foerster**

***Psyttoma latilabris* (Chen & Weng, 2005), comb. n.**

http://species-id.net/wiki/Psyttoma_latilabris

Figures 1–13

Opius latilabris Chen & Weng, 2005: 110 (Chinese key), 112–113 (description in Chinese; fig. 47), 180 (English key), 199 (description in English) (examined).


Type material. Holotype (Beneficial Insects Laboratory, Fujian Agriculture and Forestry University, Fuzhou), ♀, “[China:], Hubei, Shennongjia, Honghua, 2.vi.1988, Zhang Li-qin”. Paratype: 1 ♂, “[China:], Hubei, Shennongjia, Yangriwan, 20.vi.1988, Yang Jian-quan”. The holotype has the head missing and the wing venation is bleached.

Additional material. 1 ♀ (RMNH), “N. China: Shandong, Anqiu, Suotou Mt., 31.vii.2009, c. 120 m, Li Xi-Ying, RMNH’09”, “CVA4245, sp. 11”; 1 ♂ (RMNH), same label data.

Diagnosis. Clypeus narrow, 5 times as wide as high and 0.7 times as wide as face and face distinctly transverse (Fig. 6); pronope rather large and round; medio-posterior depression of mesoscutum medium-sized and round (Fig. 5); precoxal sulcus slightly impressed and smooth; scutellum rugulose medio-posteriorly; body completely brownish-yellow (Fig. 1).

Description. Described from ♀ collected in Anqiu, length of body 2.0 mm, of fore wing 2.1 mm.

Head. Antenna with 25 segments and 1.2 times as long as fore wing; third segment 1.1 times as long as fourth segment, length of third, fourth and penultimate segments 3.5, 3.2 and 2.7 times their width, respectively; length of maxillary palp 0.9 times height of head; labial palp segments slender; labrum slanted backwards, leaving a large space below clypeus (Fig. 9); occipital carina widely removed from hypostomal carina and dorsally absent; hypostomal carina narrow; length of eye in dorsal view 2.1 times temple; frons slightly depressed behind antennal sockets, medially convex and glabrous, smooth; face smooth, medially weakly elevated; width of clypeus 5.0 times its maximum height and 0.7 times width of face; clypeus weakly convex, distinctly protruding forwards, punctate and its ventral margin thick and slightly concave; hypoclypeal depression wide and deep (Figs 6, 9); malar suture present; without punctures between malar suture and clypeus; mandible somewhat constricted medially and gradually widened baso-ventrally, with narrow ventral carina (Fig. 9), second tooth minute.



Figures 2–11. *Psyttoma latilabris* (Chen & Weng), female from Anqiu. **2** wings **3** mesosoma dorsal **4** propodeum and first-third metasomal tergites dorsal **5** mesoscutum and scutellum dorso-lateral **6** head anterior **7** ovipositor, sheath and hypopygium lateral **8** pronotum dorsal **9** mandible lateral **10** antenna **11** hind leg.

Mesosoma. Length of mesosoma 1.2 times its height; dorsal pronope large, round and pronotum oblique anteriorly (Fig. 8); pronotal sides smooth but oblique groove crenulate and posterior groove largely absent; epicnemial area smooth dorsally; precoxal sulcus medially superficially impressed, smooth as rest of mesopleuron; pleural sulcus smooth; mesosternal sulcus deep and narrow and very finely crenulate; notauli absent on disk, only anteriorly with pair of short smooth impressions; mesoscutum glabrous and strongly shiny; medio-posterior depression of mesoscutum medium-sized, deep, round; scutellar sulcus narrow and finely crenulate laterally, widened medially; scutellum convex medially, depressed and rugulose medio-posteriorly; surface of propodeum smooth, except for superficial rugulae posteriorly (Fig. 4).

Wings. Fore wing: pterostigma elongate triangular (Fig. 2); 1-R1 ending before wing apex and 1.4 times as long as pterostigma (Fig. 2); $r:3-SR:SR1 = 3:27:55$; $2-SR:3-SR:r-m = 15:27:7$; r slender; 1-M slightly curved and SR1 straight; m-cu postfurcal; cu-a postfurcal and 1-CU1 widened; first subdiscal cell closed, CU1b short; apical third of M+CU1 sclerotized. Hind wing: M+CU:1-M:1r-m = 15:17:6; cu-a straight; m-cu present as faintly pigmented trace.

Legs. Length of femur, tibia and basitarsus of hind leg 2.3, 7.0 and 4.3 times as long as wide, respectively (Fig. 11); hind femur and tibia with medium-sized setae.

Metasoma. Length of first tergite equal to its apical width, its surface evenly moderately convex and rather densely longitudinally rugulose and dorsal carinae developed in basal 0.4 of tergite, straight (Fig. 4); second suture absent; second and following tergites smooth; length of setose part of ovipositor sheath 0.29 times fore wing, 2.7 times first tergite and equal to length of hind tibia; hypopygium distinctly acute apically and about 0.3 times as long as metasoma.

Colour. Brownish-yellow; antenna (but scapus yellowish), stemmaticum, ovipositor sheath, pterostigma and veins dark brown; palpi, mandible, tegulae and legs (but telotarsi darkened) pale yellow; wing membrane subhyaline.


Variation. Male from Anqiu has length of fore wing 2.0 mm, antenna with 24 segments, hind femur 3.0 times as long as wide and medio-anterior veins of hind wing strongly widened (Fig. 13).

Molecular data. 16S and 28S (CVA4245); GenBank Accession numbers JQ736254 and JQ736282, respectively.

Distribution. China (Shandong, Hubei).

Biology. Unknown.

Notes. In the key by Chen and Weng (2005) to the Opiinae of China the type species runs to *Phaeditoma louiseae* (Weng & Chen, 2005) comb. n., but this species has the propodeum with a pentagonal areola, the mesosoma brown, the first tergite 2.3 times longer than its apical width, the hind femur 6.0 times longer than wide, the antenna 1.6 times longer than body and the clypeus 1.7 times wider than high. Only by examining all Opiinae types in Beneficial Insects Lab., Fujian Agriculture and Forestry University (Fuzhou) it was found by the second author that the headless holotype of *Opius latilabris* Chen & Weng, 2005, from Hubei (Oriental China) is conspecific with the specimens from the Palearctic Anqiu.



Figures 12–13. *Psyttoma latilabris* (Chen & Weng), female from Anqiu, but 13 male from Anqiu, hind wing. Scale-line: **12** 1.0 ×, **13** 1.1 ×.

In the key by Tobias (1998) to the East Palaearctic Opiinae it runs to *Opius wachsmanni* Szépligeti, 1898, described from Hungary (holotype examined by the second author). This species belongs to the same genus and can be separated as follows:

- 1 Eye in dorsal view about twice as long as temple; vein CU1b of fore wing much shorter than vein 3-CU1 (Fig. 1); first tergite rather densely longitudinally rugulose (Fig.); vein 1-R1 (= metacarp) about 1.4 times as long as pterostigma (Fig. 2); head dorsally, mesosternum, mesoscutal lobes medially, metanotum, propodeum, third and following metasomal tergites brownish-yellow (Fig. 1); East Palaearctic and North Oriental.....
..... ***P. latilabris* (Chen & Weng, 2005) comb. n.**
- Eye in dorsal view about as long as temple; vein CU1b of fore wing somewhat shorter than vein 3-CU1; first tergite mainly coriaceous-punctate; vein 1-R1 slightly longer than pterostigma; head dorsally, mesosternum, mesoscutal lobes medially, metanotum, propodeum, third and following metasomal tergites black; West Palaearctic***P. wachsmanni* (Szépligeti, 1898) comb. n.**

The specimens reported as *Opius wachsmanni* Szépligeti from the East Palaearctic region (Central Asia up to Korea) by Papp (1981), Tobias and Jakimavicius (1986) and Tobias (1998) need to be re-examined and most likely belong to *P. latilabris* (Chen & Weng, 2005).

Acknowledgements

The second author wishes to thank Prof. Dr Jia-hua Chen and Mr Min-lin Zheng, Beneficial Insects Lab., Fujian Agriculture and Forestry University, Fuzhou, Fujian, for their hospitality and for the possibility to examine the Opiine types.

References

- van Achterberg C (1988) Revision of the subfamily Blacinae Foerster (Hymenoptera, Braconidae). Zoologische Verhandelingen Leiden 249: 1–324.
- van Achterberg C (1990) Illustrated key to the subfamilies of the Holarctic Braconidae (Hymenoptera: Ichneumonoidea). Zoologische Mededelingen Leiden 64: 1–20.
- van Achterberg C (1993) Illustrated key to the subfamilies of the Braconidae (Hymenoptera: Ichneumonoidea). Zoologische Verhandelingen Leiden 283: 1–189.
- van Achterberg C (1997a) Braconidae. An illustrated key to all subfamilies. ETI World Biodiversity Database CD-ROM Series Amsterdam.
- van Achterberg C (1997b) Revision of the Haliday collection of Braconidae (Hymenoptera). Zoologische Verhandelingen Leiden 314: 1–115.
- van Achterberg C, Salvo A (1997) Reared Opiinae (Hymenoptera: Braconidae) from Argentina. Zoologische Mededelingen Leiden 71(18): 189–214.
- van Achterberg C (2004a) New Indo-Australian subgenera and species of the genera *Xynobius* Foerster and *Ademoneuron* Fischer (Hymenoptera: Braconidae: Opiinae). Zoologische Mededelingen Leiden 78(20): 313–329.
- van Achterberg C (2004b) *Bitomoides* gen. nov. (Hymenoptera: Braconidae: Opiinae) from Europe. Zoologische Mededelingen Leiden 78(21): 331–335.
- van Achterberg C, Chen X-X (2004) Six new genera of Braconidae (Hymenoptera) from China. Zoologische Mededelingen Leiden 78(2): 77–100.
- Chen J-H, Weng R-Q (2005) Systematic studies on Opiinae of China (Hymenoptera: Braconidae). Fujian Science and Technology Publishing House, Fujian, volume I-III, 1–2, 1–9, 1–269.
- Fischer M (1972) Hymenoptera Braconidae (Opiinae I). (Paläarktische Region). Das Tierreich 91: 1–620.
- Fischer M (1977) Hymenoptera Braconidae (Opiinae II). (Amerika). Das Tierreich 96: 1–1001.
- Fischer M (1986) Neue Bestimmungsschlüssel für paläarktische Opiinae, neue Subgenera, Redeskriptionen und eine neue Art (Hymenoptera, Braconidae). Annalen des Naturhistorischen Museums in Wien 88/89: 607–662.
- Fischer M (1987) Hymenoptera Braconidae (Opiinae III) - äthiopische, orientalische, australische und ozeanische Region. Das Tierreich 104: 1–734.
- Gimeno C, Belshaw RD, Quicke DLJ (1997) Phylogenetic relationships of the Alysiinae/ Opiinae (Hymenoptera: Braconidae) and the utility of cytochrome b, 16S and 28S D2 rRNA. Insect Molecular Biology 6(3): 273–284. doi: 10.1046/j.1365-2583.1997.00181.x

- Li X-Y, Achterberg C van, Tan J-C (in prep.) Revision of the subfamily Opiinae (Hymenoptera: Braconidae) from Hunan (China), with the description of thirty-six new species and one new genus.
- Papp J (1981) Braconidae (Hymenoptera) from Korea V. *Acta Zoologica Academiae Scientiarum Hungaricae* 27: 139–158.
- Tobias VI, Jakimavicius A (1986) Alysiinae and Opiinae. In: Medvedev GS (Ed.) *Opredelitel Nasekomykh Evropeiskoi Tsasti SSSR* 3, *Peredpontdatokrylye* 4. *Opr. Faune SSSR* 147 (3)5, 7–231.
- Tobias VI (1998) Opiinae. In: Ler PA (Ed.) *Key to the insects of Russian Far East*. 4. *Neuropteroidea, Mecoptera, Hymenoptera* 3. *Dal'nauka, Vladivostok*, 558–655.
- Wharton RA (1988) Classification of the Braconid subfamily Opiinae (Hymenoptera). *Canadian Entomologist* 120: 333–360. doi: 10.4039/Ent120333-4
- Wharton RA (1997) Generic relationships of opiine Braconidae (Hymenoptera) parasitic on fruit-infesting Tephritidae (Diptera). *Contributions of the American Entomological Institute* 30(3): 1–53.
- Yu DS, Achterberg K van, Horstmann K (2009) Biological and taxonomical information: Ichneumonoidea 2006. *Taxapad Interactive Catalogue*, Lexington.