

First record of the parasitoid *Gonatopus flavipes* Olmi, 1984 (Hymenoptera, Dryinidae) in Brazil's Amazon forest

André Luis Martins¹, Diones Krinski^{2,3}

1 Programa de Pós-Graduação em Entomologia, Departamento de Zoologia, Laboratório de Biologia Comparada de Hymenoptera, Universidade Federal do Paraná, CEP 81531-980, Curitiba, PR, Brazil **2** Programa de Pós-Graduação em Zoologia, Departamento de Zoologia, Laboratório de Controle Integrado de Insetos, Universidade Federal do Paraná, CEP 81351-980 Curitiba, PR, Brazil **3** Departamento de Ciências Biológicas, Universidade do Estado de Mato Grosso, CEP 78300-000 Tangará da Serra, MT, Brazil

Corresponding author: Andre L. Martins (andrelmartinsbiol@hotmail.com)

Academic editor: Michael Ohl | Received 18 April 2016 | Accepted 8 May 2016 | Published 27 June 2016

<http://zoobank.org/3CBFB7D1-2735-47A3-8B87-BDCE8D03642F>

Citation: Martins AL, Krinski D (2016) First record of the parasitoid *Gonatopus flavipes* Olmi, 1984 (Hymenoptera, Dryinidae) in Brazil's Amazon forest. Journal of Hymenoptera Research 50: 191–196. doi: 10.3897/JHR.50.8897

Abstract

This study reports for the first time the occurrence of *Gonatopus flavipes* Olmi, 1984 in Pará State, Brazil. Specimens were collected on upland rice crops of Cambará variety in Novo Progresso (7°07'45.71"S 55°23'21.13"W). A sampling of insects with pitfall traps was conducted between November 2010 and March 2011. Specimens of *G. flavipes* were identified, illustrated and deposited in the Entomological Collection of the Department of Zoology, Federal University of Paraná, Curitiba, Brazil (DZUP/UFPR). This record indicates that rice crops may represent important habitats for this species.

Keywords

Chrysidoidea, Gonatopodinae, *Gonatopus* group seven, pitfall traps

Introduction

Gonatopus Ljung, 1810 is one of the 11 genera belonging to the subfamily Gonatopodinae (Hymenoptera, Dryinidae). With about 440 worldwide described species, among which 120 known in the Neotropics and 32 in Brazil (Olmí and Virla 2014; Martins et al. 2015a, b), this genus is divided into 12 groups (nine registered for the Neotropics and five for Brazil) (Olmí and Virla 2014). Little is known about the biology of parasitoids belonging to the genus *Gonatopus*. The few studies report the association of the genus with some families of leafhoppers and planthoppers (Hemiptera, Auchenorrhyncha): Acaloniidae, Caliscelidae, Cicadellidae, Cixiidae, Delphacidae, Dictyopharidae, Flatidae, Issidae, Lophopidae, Meenoplidae and Tropiduchidae (Guglielmino et al. 2013; Olmí and Virla 2014).

Gonatopus flavipes Olmí, 1984, broadly distributed from Mexico to Argentina, belongs to group seven, which is the largest group of *Gonatopus* with exactly 61 described species. Both sexes of *G. flavipes* are known (Olmí and Virla 2014). The following species of Cicadellidae (Hemiptera, Auchenorrhyncha) are reported as hosts: *Mendozellus asuntia* Cheng in Argentina; *Dalbulus maidis* (DeLong and Wolcott) in Piauí, Brazil; *Dalbulus elimatus* (Ball), *Planicephalus flavicosta* (Stål) and *Graminella comata* (Ball) in Mexico (Guglielmino and Olmí 1997, 2006; Meneses et al. 2013; Moya-Raygoza 1990, 1993; Olmí and Virla 2014; Virla 1992).

The Companhia Nacional de Abastecimento do Brasil (CONAB 2016) records rice cultivation in Pará State since 1976–1977, mainly in the North, with recent planting of upland rice in the Southwest (less than 10 years) (Azevedo 2009; Krinski 2014; Lopes et al. 2004; Silva and Magalhães 1981). For this reason, a few entomological studies have been conducted on rice cultivars of this region. This study reports for the first time the occurrence of *G. flavipes* in upland rice crop in Southwestern region of Pará State (Brazil).

Materials and methods

A sampling of insects was conducted with pitfall traps on upland rice crops (Cambará variety), in Novo Progresso, State of Pará, Brazil (7°07'45.71"S 55°23'21.13"W) (Fig. 1), between November 2010 and March 2011. The sampling effort included 16 collecting points visited weekly, covering 400 meters in four transects of 100 meters, distant 25 meters from each other.

The collected specimens of *Gonatopus* were sent to the Laboratory of Comparative Hymenoptera Biology at Federal University of Paraná (UFPR), where they were identified to species level using a stereo-microscope LEICA M125 coupled to digital camera LEICA DFC295. The images were processed by Zerene Stacker software (1.04 version build). Digital scanning electronmicroscope (SEM) photographs were taken with a TESCAN VEGA3 LMU in low vacuum mode. The figures were prepared using Adobe Photoshop (version 11.0). Specimens are deposited in the Entomological Collection of the Department of Zoology, Federal University of Paraná (DZUP).




Figure 1. Distribution map of *Gonatopus flavipes*. Black circle: record areas in Argentina, Bolivia, Brazil (Piauí, São Paulo and Santa Catarina), Costa Rica, Ecuador, Guadeloupe, Jamaica and Mexico; Star: new record of occurrence (Florentino farm – municipality of Novo Progresso, Pará State, Brazil). Source: i3Geo (free software). Licensed as General Public License (GNU) and created by Ministry of Environment (MMA).

Results and discussion

Gonatopus flavipes Olmi, 1984

Gonatopus flavipes Olmi, 1984: 1752.


Note. Only two female specimens were collected (Figs 2–5). They present the following diagnostic features:

Diagnosis. Completely yellow testaceous except petiole black and metasoma partly brown (Fig. 2). Pronotum shiny, unsculptured, crossed by strong transverse furrow. Scutum shiny, unsculptured, laterally with two pointed apophyses (Fig. 3). Meso-metapleural suture obsolete (Fig. 2). Head with frontal line complete; occipital carina absent; OL (distance between the inner edges of a lateral ocellus and the median ocellus)/POL (distance between the inner edges of the lateral ocelli) = 1.5/1.0 (Fig. 4). Enlarged claw with one small subapical tooth and one row of six peg-like hairs. Segment 5 of protarsus with two rows of 22 + 12 lamellae (Fig. 5). Tibial spurs 1/0/1.

Material examined. Two females: Brazil, PA, Novo Progresso, Florentino Farm, Upland rice crop, 7°08'41"S 55°22'43"W, 09.ii.2011, D. Krisnki, Pitfall traps (DZUP).

Gonatopus flavipes was recorded for Argentina, Bolivia, Brazil (Piauí, São Paulo and Santa Catarina States), Costa Rica, Ecuador, Guadeloupe, Jamaica and Mexico (Olmi and Virla 2014) (Fig. 1). In addition we record for the first time the presence of this species from Pará State, Brazil, in upland rice crop.

Therefore, considering the economic importance of rice crops in Brazil, we recommend careful monitoring in rice areas and studies on the biology, morphology and ecology of *G. flavipes* in different rice crops.



Figures 2–5. *Gonatopus flavipes*. **2** Habitus **3** Mesosoma in dorsal view **4** Head in dorsal view **5** Chela.

In addition, more studies are needed to assess the population fluctuation of this parasitoid in different rice varieties, mainly to investigate if this species can be used in the future for the biological control of Auchenorrhyncha (Guglielmino 2002; Mita et al. 2012). In fact, research out of Brazil have reported the occurrence of *Gonatopus* species associated with important pests of rice (Mita and Pham 2014).

Acknowledgements

Authors acknowledge the farmers of Fazenda Florentino, Marlete Florentino, Eurides Florentino (*in memoriam*), and Nadir de Lima Florentino (*in memoriam*) for allowing this research on their property. We also thank the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) and the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES) for fellowships provided to D. Krinski (CNPq – Proc.: 141243/2012-0 and CAPES – Proc.: 939980) and A. L. Martins (CNPq – Proc.: 142415/2015-4). Many thanks to the Center of Electron Microscopy of Universidade Federal do Paraná (UFPR) for the SEM photomicrographs. Many thanks to Dr. Massimo Olmi (Tropical Entomology Research Center, Viterbo, Italy) for help in the identification of species and for the review of the manuscript.

References

- Azevedo R de (2009) Manejo integrado de insetos-praga em arroz, milho e soja. Simpósio sobre Manejo de Pragas, 145–160. <http://ainfo.cnptia.embrapa.br/digital/bitstream/item/46112/1/Azevedo00186.pdf> [accessed 05 Apr 2016]
- CONAB (2016) CONAB (Companhia Nacional de Abastecimento), 2014. Arroz. Séries Históricas de Área Plantada, Produtividade e Produção, Relativas às Safras 1976/77 a 2014/15. <http://www.conab.gov.br/conteudos.php?a=1252&t=2> [accessed 05 Apr 2016]
- Guglielmino A, Olmi M (1997) A host-parasite catalog of world Dryinidae (Hymenoptera: Chrysidoidea). Contributions on Entomology International 2(2): 165–298.
- Guglielmino A (2002) Dryinidae (Hymenoptera Chrysidoidea): an interesting group among the natural enemies of the Auchenorrhyncha (Hemiptera). Denisia 4: 549–556.
- Guglielmino A, Olmi M (2006) A host-parasite catalog of world Dryinidae (Hymenoptera: Chrysidoidea): first supplement. Zootaxa 1139: 35–62.
- Guglielmino A, Olmi M, Bückle C (2013) An updated host-parasite catalogue of world Dryinidae (Hymenoptera: Chrysidoidea). Zootaxa 3740(1): 001–113. doi: 10.11646/zootaxa.3740.1.1
- Krinski D (2014) Artrópodes associados ao arroz de terras altas, *Oryza sativa*, em Novo Progresso, estado do Pará: níveis de danos e estratégias para manejo. Universidade Federal do Paraná, Paraná, 326 pp. [Tese de doutorado em Ciências – Zoologia]
- Ljungh SJ (1810) *Gonatopus*, novum insectorum genus. Beiträge zur Naturkunde 2: 161–163.
- Lopes AM, Silveira-Filho A, Husny JCE, Correa JRV (2004) BRS Talento: cultivar de arroz para cultivo em terra firme no Estado do Pará. Embrapa Amazônia Oriental – Comunicado Técnico 93. <http://ainfo.cnptia.embrapa.br/digital/bitstream/item/18749/1/com.tec.93.pdf> [accessed 05 Apr 2016]
- Martins AL, Lara RIR, Periotto NW (2015a) New records of Dryinidae (Hymenoptera: Chrysidoidea) from the Atlantic Rainforest of São Paulo, Brazil. The Pan-Pacific Entomologist 91(2): 196–199. doi: 10.3956/2015-91.2.196.
- Martins AL, Lara RIR, Periotto NW, Olmi M (2015b) Two new species of Dryinidae (Hymenoptera: Chrysidoidea) from areas of Atlantic Rainforest at São Paulo State, Brazil. Brazilian Journal of Biology 75(2): 455–459. doi: 10.1590/1519-6984.19613
- Meneses AR, Queiroz RB, Olmi M, De Oliveira CM, Silva PRR, Bezerra VS (2013) Descoberta de *Gonatopus flavipes* (Olmi) como um novo parasitóide de *Dalbulus maidis* (De-Long & Wolcott). In: Anais do 13º SICONBIOL, Simpósio de Controle Biológico, 15–18 Setembro 2013, Bonito, Mato Grosso do Sul, Brazil.
- Mita T, Pham HT (2014) Dryinidae collected from rice paddy in Vietnam (Hymenoptera: Chrysidoidea). Japanese Journal of Systematic Entomology 20(2): 245–246.
- Mita T, Matsumoto Y, Sanada-Morimura S, Matsumura M (2012) Passive long Distance migration of apterous dryinid wasps parasitizing rice planthoppers. In: Stevens L (Ed.) Global Advances in Biogeography. InTech, Rijeka, 49–60. <http://www.intechopen.com/books/global-advances-in-biogeography/passive-long-distance-migration-ofapterous-dryinid-wasps-using-rice-planthopper>

- Moya-Raygoza G (1990) Parasitoides de *Dalbulus* spp. (Homoptera: Cicadellidae) en Jalisco, México. Tesis de maestría en ciencias, Colegio de Postgraduados, Chapingo, México, 62 pp.
- Moya-Raygoza G (1993) Diversity of leafhoppers and their hymenopteran parasitoids in maize, teosinte and gamagrass. Proceedings of 8th Auchenorrhyncha Congress, Delphi, Greece, 9–13 August 1993, 67–68.
- Oلمي M (1984) A revision of the Dryinidae (Hymenoptera). Memoirs of the American Entomological Institute, Gainesville 37: 1–1913.
- Oلمي M, Virla EG (2014) Dryinidae of the Neotropical region (Hymenoptera: Chrysidoidea). Zootaxa 3792(1): 1–534. doi: 10.11646/zootaxa3792.2.1
- Silva AB, Magalhães BP (1981) Insetos novíços à cultura do arroz no Estado do Pará. Embrapa Amazônia Oriental – Circular Técnica 22. <https://www.infoteca.cnptia.embrapa.br/bitstream/doc/379480/1/CPATUCirTec22.pdf> [accessed 05 Apr 2016]
- Virla EG (1992) Estudio bionómico de parasitoides e hiperparasitoides de Homópteros Cica-deloideos Argentinos. Trabajo de tesis para optar al título de Doctor en Ciencias Naturales, Facultad de Ciencias Naturales y Museo, Universidad de La Plata, Argentina, 263 pp.